


HPLC Method for Analysis of Kinetin Phytohormone on Primesep 100 Column



Separation type: Liquid Chromatography Mixed-mode SIELC Technologies

Kinetin is a type of cytokinin, a class of phytohormones that play a crucial role in promoting cell division, growth, and delaying aging (senescence) in plants. It was one of the first cytokinins discovered and is often used in plant tissue culture and agriculture to enhance plant growth and development. Kinetin is a powerful cytokinin that is critical for cell division, shoot formation, and delaying aging in plants. Its application spans plant tissue culture, agriculture, and even the cosmetic industry.

Kinetin was first isolated from degraded DNA in herring sperm in 1955, though it is not commonly found in high concentrations in plants compared to other cytokinins like zeatin. However, kinetin is naturally occurring and has been detected in various plant tissues.

Kinetin binds to cytokinin receptors in plants, initiating a signaling cascade that leads to gene expression changes, promoting cell division and growth. Its action is often in opposition to auxins, and both hormones work together to balance root and shoot growth.

Kinetin can be retained, separated and analyzed using a Primesep 100 mixed-mode stationary phase column. The analysis employs an isocratic method with a simple mobile phase comprising water, acetonitrile (MeCN), and sulfuric acid as a buffer. This method allows for detection using UV 210 nm.

You can find detailed UV spectra of Kinetin and information about its various lambda maxima by visiting the following link .

Method Parameters

Column	Primesep 100, 4.6 x 150 mm, 5 µm, 100 Å, dual ended
Mobile Phase	MeCN – 40%
Buffer	H2SO4 -0.2%
Flow Rate	1.0 mL/min
Detection	UV 210 nm
Sample	0.1 mg/ml

Quelle: <https://sielc.com/hplc-method-kinetin>