


## HPLC-MS Method for Analysis of Oxytocin on Primesep 200 Column



<b>Column:</b>	Primesep 200
<b>Column size:</b>	2.1 × 100 mm, 5 µm
<b>Column part number:</b>	200-21.100.0510
<b>Mobile phase:</b>	MeCN/H <sub>2</sub> O – 30/70%
<b>Buffer:</b>	Ammonium format pH 3.0 – 20 mM
<b>Flow rate:</b>	0.2 mL/min
<b>Detection:</b>	UV 275 nm, SIM 1008+

Separation type: Liquid Chromatography Mixed-mode

Oxytocin is a neuropeptide and a hormone that plays a significant role in social bonding, sexual reproduction, childbirth, and the period after childbirth.

Oxytocin is a peptide hormone and neuropeptide that plays a crucial role in social bonding, reproduction, and childbirth. It is often called the “love hormone” or “bonding hormone” because of its association with social interactions, including maternal behaviors, bonding between couples, and even the bond between humans and their pets.

Oxytocin is a neuropeptide and a hormone that plays a significant role in social bonding, sexual reproduction, childbirth, and the period after childbirth. Here’s a breakdown of its classification and functions:

1. Chemical Classification:

2. Production and Release:

Oxytocin can be retained, and analyzed on a Primesep 200 mixed-mode stationary phase column using an isocratic analytical method with a simple mobile phase of water, Acetonitrile (MeCN), and a Ammonium format as a buffer. This analysis method can be detected in the UV 275 nm.

## Method Parameters

<b>Column</b>	Primesep 200, 2.1 x 100 mm, 5 µm, 100 Å, dual ended
<b>Mobile Phase</b>	MeCN/H <sub>2</sub> O – 30/70%
<b>Buffer</b>	Ammonium Format pH 3.0- 20 mM
<b>Flow Rate</b>	0.2 mL/min
<b>Detection</b>	UV 275 nm, LCMSSIM1008+

Quelle: <https://sielc.com/hplc-determination-of-oxytocin>