


HPLC Separation of Metribuzin on Mixed-Mode HPLC Columns



Separation type: Liquid Chromatography Mixed-mode

High Performance Liquid Chromatography (HPLC) Method for Analysis of Metribuzin. Metribuzin is a pre- and post- emergence herbicide on the EURL (European Union Reference Laboratory) Target Pesticide List for the EUPT-CF9 2015 (European Union Proficiency Test for Cereals and Feeding stuff). Metribuzin disrupts photosynthesis in plants growing near food crops such as potatoes, tomatoes and sugar cane. Obelisc R and Primesep 100 were used to retain Metribuzin using multiple modes of separation. Obelisc R contains a long hydrophobic chain and ion-pairing groups, while Primesep 100 contains embedded acidic ion-pairing groups. Method is LC/MS compatible and can be used with similar conditions to retain many pesticides.

SIELC has developed the Obelisc™ columns, which are mixed-mode and utilize Liquid Separation Cell technology (LiSC™). These cost-effective columns are the first of their kind to be commercially available and can replace multiple HPLC columns, including reversed-phase (RP), AQ-type reversed-phase, polar-embedded group RP columns, normal-phase, cation-exchange, anion-exchange, ion-exclusion, and HILIC (Hydrophilic Interaction Liquid Chromatography) columns. By controlling just three orthogonal method parameters - buffer concentration, buffer pH, and organic modifier concentration - users can adjust the column properties with pinpoint precision to separate complex mixtures.

Method Parameters

| | |
|---------------------|--|
| Column | Primesep 100, 2.1×150 mm, 5 µm, 100 Å |
| Mobile Phase | Gradient MeCN – 10-70%, 15 min |
| Buffer | Gradient AmAc pH 3.0- 20-60 mM, 15 min |
| Flow Rate | 0.4 mL/min |
| Detection | UV, 270 nm |

Quelle: <https://sielc.com/Application-HPLC-Separation-of-Metribuzin-on-Mixed-Mode-HPLC-Columns>