


HPLC Method for Analysis of Crystal Violet on Primesep 100 Column

<https://sielc.com/hplc-determination-of-crystal-violet>

Chromatogram



Column:	Primesep 100
Column size:	4.6 × 150 mm, 5 μm
Column part number:	100-46.150.0510
Mobile phase:	MeCN/H ₂ O – 80/20%
Buffer:	H ₂ SO ₄ - 0.2%
Flow rate:	1.0 mL/min
Detection:	Vis 590 nm

Description

· Separation type: Liquid Chromatography Mixed-mode · HPLC Method for Analysis of Crystal Violet on Primesep 100 Column by SIELC Technologies

Crystal Violet, also known as Methyl Violet 10B, is a synthetic dye belonging to the triarylmethane dye family. It has the molecular formula C₂₅H₃₀N₃. This dye is primarily known for its use in microbiology as a stain, but it has other applications as

well.

Crystal Violet can be retained, and analyzed on a Primesep 100 mixed-mode stationary phase column using an isocratic analytical method with a simple mobile phase of water, Acetonitrile (MeCN), and a sulfuric acid (H₂SO₄) buffer. This analysis method can be detected in the UV-Vis regime at 540, 590, and 200 nm.

Method Parameters

Mobile Phase	MeCN/H ₂ O – 80/20%
Buffer	H ₃ PO ₄ – 0.2%
Flow Rate	1.0 ml/min
Detection	Vis, 590 nm
Peak Retention Time	9.37 min
Sample concentration	0.0006 mg/ml
Injection volume	5 µl
LOD	10 ppb
Class of Compounds	Dyes
Analyzing Compounds	Crystal Violet

HPLC Column Used

Primesep 100, 4.6 x 150 mm, 5 µm, 100 A, dual ended

[Order this column at hplc-shop.de](http://hplc-shop.de) →