



HPLC Determination of Shikimic Acid on Newcrom BH Column

<https://sielc.com/hplc-determination-of-shikimic-acid>

Chromatogram

Column: Newcrom BH
Column size: 4.6 × 150 mm, 5 µm
Mobile phase: MeCN/H₂O – 5/95%
Buffer: H₃PO₄ – 0.02%
Flow rate: 1 ml/min
Detection: UV 200 nm

Shikimic acid



Description

· High Performance Liquid Chromatography (HPLC) Method for Analysis of Shikimic Acid

Shikimic Acid, also known as shikimate, is a well-known biochemical intermediary integral to the metabolic process found in plants and microorganisms. It was first isolated from the Japanese shikimi flower, the Japanese star anise, which is where it gets its name. It has the chemical formula C₇H₁₀O₅. This cyclohexanecarboxylic acid can be detected in the low UV regime. Using a Newcrom BH mixed-mode column and a mobile phase consisting of (mostly) water and acetonitrile, shikimic acid can be retained, measured, and analyzed. This analysis method can be UV detected at 200 nm with high resolution.

Method Parameters

Mobile Phase	MeCN/H ₂ O – 5/95%
Buffer	H ₃ PO ₄ – 0.02%
Flow Rate	1.0 ml/min
Detection	UV 200 nm

Class of Compounds	Acid
Analyzing Compounds	Shikimic acid

HPLC Column Used

Newcrom BH, 4.6 x 150 mm, 5 µm, 100 A, dual ended

[Order this column at hplc-shop.de →](https://www.hplc-shop.de)