


HPLC Method for Analysis of 3'-Deoxyadenosine and 2'-Deoxyadenosine on BIST B+ Column

<https://sielc.com/hplc-method-of-deoxy>

Chromatogram



Column:	BIST B+
Column size:	4.6 × 150 mm, 5 μm
Column part number:	TBP-46.150.0510
Mobile phase:	MeCN – 85%
Buffer:	H ₃ PO ₄ - 0.2%
Flow rate:	1.0 mL/min
Detection:	UV 260 nm

Description

HPLC Method for Analysis of Cordycepin , Deoxyadenosine on Cordycepin , Deoxyadenosine by SIELC Technologies .

Cordycepin is an adenosine analogue with the chemical formula $C_{10}H_{13}N_5O_3$. It is reported to prevent cell reproduction in various cancer cells. It also might possess antioxidant and anti-inflammatory properties, when considered in addition with its ability to cross the blood-brain barrier, it may become widely used in pharmaceuticals.

Deoxyadenosine is a deoxyribonucleoside with the chemical formula $C_{10}H_{13}N_5O_3$. It is a derivative of adenosine. High presence of it can kill T lymphocytes and kill those cells, leading to adenosine deaminase severe combined immunodeficiency disease, also known as ADA-SCID.

Cordycepin, Deoxyadenosine can be retained and analyzed using the BIST B+ stationary phase column. The analysis utilizes an isocratic method with a simple mobile phase consisting of water and acetonitrile (MeCN) with a Phosphoric Acid buffer. Detection is performed using UV.

Method Parameters

Mobile Phase	MeCN – 85%
Buffer	H ₃ PO ₄ – 0.2%
Flow Rate	1.0 ml/min
Detection	UV 260 nm
Peak Retention Time	8.2, 9.8 min
Class of Compounds	Nucleosides
Analyzing Compounds	Cordycepin, Deoxyadenosine

HPLC Column Used

BIST B+, 4.6 x 150 mm, 5 µm, 100 Å, dual ended

[Order this column at hplc-shop.de](http://hplc-shop.de) →